

Notation for general quantum teleportation

Chong Li¹, He-Shan Song and Yao-Xing Luo

*Department of Physics, Dalian University of Technology
Dalian, 116024, People's Republic of China.*

Abstract

A new notation for the quantum teleportation of finite dimensional quantum state through a generally entangled quantum channel is introduced. For a given quantum channel an explicit mathematical criterion that governs the faithful teleportation is presented.

¹ lichong@student.dlut.edu.cn

Quantum teleportation has been introduced by Benett et al. [1] and discussed by a number of authors for two dimensional quantum state[2][3][4][5]. By means of a classical and distributed quantum communication channel, an unknown quantum state is destroyed at the sending place while its perfect replica state, after some rotation, appears at remote receiving place. Recently, for teleportation of a general finite dimensional quantum state has been discussed in [6] and [7], and the condition of the unitary transformation was given in the[7].

In this letter, we introduced a new notation for quantum teleportation through a generally entangled quantum channel and constructed an explicit unitary transformation U which receiver Bob performed on his qubit. It is adapt to teleport a n -dimensional quantum state by an $(n + 1)$ -dimension generally entangled channel. At last, we given all the unitary transformation U for the case that both channel and measurement are in the *Bell* state (maximally entangled state).

Let H_1 be the *Hilbert* space with dimensions N , and $\{|i\rangle_1 | i = 1 \dots N\}$ be the set of orthogonal basis in H_1 . Alice has a general state $|\varphi\rangle_1$ unknown to her in the *Hilbert* space H_1 as,

$$|\varphi\rangle_1 = \sum_{i=1}^N \alpha_i |i\rangle_1. \quad (1)$$

Where $\alpha_i \in \mathfrak{C}$, and $\sum_{i=1}^N |\alpha_i|^2 = 1$. For convenience we name an arbitrary dimensional quantum state as a qubit.

Now we definite some useful notations:

Definition 1 $\alpha = (\alpha_1 \ \cdots \ \alpha_{N-1} \ \alpha_N)$,

Definition 2 $\mathbf{e}_i = (|1\rangle_i \ \cdots \ |N-1\rangle_i \ |N\rangle_i)$,

Definition 3 $\tilde{\mathbf{e}}_i = ({}_i\langle 1| \cdots {}_i\langle N-1| \quad {}_i\langle N|)$.

Where the states $|1\rangle_i \cdots |N\rangle_i$ represent the different states of the i -th qubit. For example, when $N = 2$, they can be spin- $\frac{1}{2}$ states, namely $|1\rangle \rightarrow |\uparrow\rangle$ and $|2\rangle \rightarrow |\downarrow\rangle$, and so on.

We rewrite the Eq.(1) by the notation as

$$|\varphi\rangle_1 = \boldsymbol{\alpha} \mathbf{e}_1^t, \quad (2)$$

where $\mathbf{e}_i^t$ denotes transpose of $\mathbf{e}_i$.

Let H_2 and H_3 be auxiliary *Hilbert* spaces attached respectively to Alice and Bob, with dimensions $N_2 = N_3 = N$. A generally entangled state of two qubits in the *Hilbert* space $H_2 \otimes H_3$ is given by

$$|\phi\rangle_{23} = \sum_{i,j}^N a_{ij} |i\rangle_2 |j\rangle_3$$

with ${}_{23}\langle\phi|\phi\rangle_{23} = 1$. Both $\{|i\rangle_2 | i = 1 \dots N\}$ and $\{|i\rangle_3 | i = 1 \dots N\}$ are sets of orthogonal basis in *Hilbert* spaces H_2 and H_3 respectively. We express $|\phi\rangle_{23}$ by the notation

$$|\phi\rangle_{23} = \mathbf{e}_2 A \mathbf{e}_3^t, \quad (3)$$

where the known matrix $A = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1N} \\ a_{21} & a_{22} & \cdots & a_{2N} \\ \vdots & a_{i2} & \cdots & \vdots \\ a_{N1} & a_{N2} & \cdots & a_{NN} \end{pmatrix}$ for given quantum channel. The state for joint system of particle 1 and 2 given to Alice and 3 given to Bob is

$$|\Psi\rangle_{123} = |\varphi\rangle_1 \otimes |\phi\rangle_{23} = \boldsymbol{\alpha} \mathbf{e}_1^t \mathbf{e}_2 A \mathbf{e}_3^t = \boldsymbol{\alpha} \mathbf{E} A \mathbf{e}_3^t \quad (4)$$

with $\mathbf{E} = \mathbf{e}_1^t \mathbf{e}_2$.

To transform the state of Bob's qubit to be $|\varphi\rangle_3$, Bob need to do an unitary transformation on his qubit, after he obtain the measured outcomes from Alice by classical channel. This process is called as quantum teleportation.

Alice make a joint measurement on the first and the second qubits, then particles 1 and 2 will form an entangled state $|\phi'\rangle_{12} = \mathbf{e}_1 B \mathbf{e}_2^t$, with $B = \begin{pmatrix} b_{11} & b_{12} & \cdots & b_{1N} \\ b_{21} & b_{22} & \cdots & b_{2N} \\ \vdots & b_{i2} & \cdots & \vdots \\ b_{N1} & b_{N2} & \cdots & b_{NN} \end{pmatrix}$. After Alice's measurement, Bob's particle 3 will have been projected into

$$|\varphi''\rangle_3 = {}_{12}\langle\phi'| \Psi\rangle_{123} = \rho \alpha X \mathbf{e}_3^t, \quad (5)$$

where $|\varphi''\rangle_3 = \rho |\varphi'\rangle_3$, $\rho X = BA$, is a known matrix and ρ is a known common factor, it is the probability amplitude of the state $|\varphi''\rangle_3$ (if the channel and measurement are *Bell* state, $\rho = \frac{1}{2}$). Bob need to perform a transformation U on his qubit in order to obtain perfect replica of $|\varphi\rangle_1$ and hence realize the teleportation.

$$U |\varphi'\rangle_3 = |\varphi\rangle_3 = \alpha \mathbf{e}_3^t \quad (6)$$

Theorem 1 If there is inverse matrix of the known matrix X^t , the quantum teleportation can be realized and $(X^t)^{-1}$ is just the transformation U that fulfills the quantum teleportation.

Proof :

By using the Eq.(2) (3) and (4), we have

$$\begin{aligned} |\varphi'\rangle_3 &= \sum_{i,j,k,m,n}^N \alpha_i a_{jk} b_{mn} \langle n|j\rangle_{21} \langle m|i\rangle_1 |k\rangle_3 \\ &= \sum_{i,j,k,m,n}^N \alpha_i a_{jk} b_{mn} \delta_{mi} \delta_{nj} |k\rangle_3 \\ &= \rho \sum_{i,k}^N \alpha_i x_{ik} |k\rangle_3 \left(\rho x_{ik} = \sum_j^N a_{jk} b_{ij} \right) \end{aligned}$$

namely

$$\left| \varphi'' \right\rangle_3 = \boldsymbol{\alpha} B A \mathbf{e}_3^t = \rho \boldsymbol{\alpha} X \mathbf{e}_3^t = \rho \left| \varphi' \right\rangle_3.$$

Bob perform the operation U on the state $\left| \varphi' \right\rangle_3$, then he will obtain perfect replica of $\left| \varphi \right\rangle_1$,

$$U \left| \varphi' \right\rangle_3 = \left| \varphi \right\rangle_3,$$

namely

$$U \sum_k^N x_{ik} \left| k \right\rangle_3 = \left| i \right\rangle_3. \quad (7)$$

Let the basis $\left| i \right\rangle$ be an N -dimensional column vector

$$\left| i \right\rangle = \begin{pmatrix} 0 & \cdots & 1 & \cdots & 0 \end{pmatrix}^t. \quad (8)$$

1..... i N

From the Eq.(7) and(8), we have

$$\sum_k^N U_{ij} X_{kj} = \delta_{ik},$$

so the matrix U is nothing but the inverse matrix of the known matrix X^t

$$U X^t = I,$$

$$U = (X^t)^{-1}. \blacksquare$$

From above discussion, we see that any finite state can be teleported through a generally entangled state.

This theorem can be extended the teleportation of multi-qubit such as the states discussed in the Ref.[8], if we re-define merely some notation. We define the state to be teleported

$$\left| \psi \right\rangle = \sum_{i=1}^N \alpha_i \left| \underbrace{ii \cdots i}_n \right\rangle_1 = \boldsymbol{\alpha} \mathbf{e}_1^t.$$

with $\mathbf{e}_1 = \left(\left| \underbrace{1 \cdots 1}_n \right\rangle_1 \left| \underbrace{2 \cdots 2}_n \right\rangle_1 \cdots \left| \underbrace{n \cdots n}_n \right\rangle_1 \right)$,
the channel is

$$|\phi\rangle_{23} = \sum_{i=1}^N a_i \left| \underbrace{ii \cdots i}_{n+1} \right\rangle_{23} = \mathbf{e}_2 A \mathbf{e}_3^t,$$

with $\mathbf{e}_2 = (|1\rangle_2 |2\rangle_2 \cdots |n\rangle_2)$ and $\mathbf{e}_3 = \left(\left| \underbrace{1 \cdots 1}_n \right\rangle_3 \left| \underbrace{2 \cdots 2}_n \right\rangle_3 \cdots \left| \underbrace{n \cdots n}_n \right\rangle_3 \right)$.
and Alice's measured result is

$$\left| \phi' \right\rangle_{12} = \mathbf{e}_1 B \mathbf{e}_2,$$

then we can construct the transformation on the third qubit by using the theorem

$$U = ((BA)^t)^{-1} / \rho.$$

It is easily show that the operation U may be non-unitary. In this case, the teleportation become probabilistic as the result given in Ref. [9].

In general, the Bob's transformation U should be unitary, so a limitation should be considered, namely the matrix X (or X^t) must satisfy the following equation,

$$(X^t)^\dagger = (X^t)^{-1} \\ \sum_{j=1}^N x_{ji} x_{kj}^* = \delta_{ik}$$

Therefore, if the channel and the Alice's measured result are known, and the matrix X is unitary, Bob can obtain the transported state. Neither the channel or the measurement result must be maximally entangled state, or the channel can be a maximally entangled state, but the Alice's measured result can be non-maximum entangled state. For example, we take $N = 2$, let the quantum channel be a generally entangled state $|\psi\rangle_{23} = \mathbf{e}_2 A \mathbf{e}_3^t$,

with $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $\det(A) \neq 0$, and let the measured result of Alice be $\left| \phi' \right\rangle_{12} = \mathbf{e}_1 B \mathbf{e}_2^t$, with $B = \begin{pmatrix} a' & b' \\ c' & d' \end{pmatrix}$ and $\det(B) \neq 0$, then the unitary matrix $(X)^t = (BA)^t / \rho = \begin{pmatrix} a'a + b'c & ac' + cd' \\ a'b + b'd & c'b + d'b \end{pmatrix} \frac{1}{\rho} = \begin{pmatrix} x_1 & x_3 \\ x_2 & x_4 \end{pmatrix}$. We can calculate out the unitary matrix X ,

$$\begin{aligned}
X &= \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ \sin(\theta) & -\cos(\theta) \end{pmatrix} \\
\text{or } X &= \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix},
\end{aligned}$$

so we can get the inverse matrix of X^t as

$$(X^t)^{-1} = (X^t)^\dagger.$$

If we assume that Alice's measurement is *Bell* basis, namely $B \in \{\sigma_0, \sigma_1, \sigma_2, \sigma_3\}$, then the matrix $(X^t)^{-1}$ can be obtained easily. We given all the matrix U in table (1) for the different channels and different measured result, and all the channels and the measured results are *Bell* state.(all the matrixes have a common factor $\rho = \frac{1}{\sqrt{2}}$)

In addition, the theorem can also be used to the general case, as $\dim(H_1) = n \leq \dim(H_2) = m$. Let α and $\mathbf{e}_1$ be respectively

$$\begin{aligned}
\alpha &= (\alpha_1 \ \cdots \ \alpha_n \ \alpha_{n+1} \ \cdots \ \alpha_m) \text{ where } \alpha_i = 0 \text{ for } i = n+1 \cdots m \\
\text{and } \mathbf{e}_1 &= (|1\rangle_1 \ \cdots \ |n\rangle_1 \ |n+1\rangle_1 \ \cdots \ |m\rangle_1).
\end{aligned}$$

If we can construct a matrix U , which made matrix $X^t U$ be diagonal and all diagonal elements of matrix $X^t U$ be 0 or 1, the teleportation can be realized. We can delete all of the elements from the $(n+1)$ -th row to the m -th row and $(n+1)$ -th column to the m -th column of the matrix X^t then we got matrix $X^{t'}$, and construct the inverse matrix of $X^{t'}$ by the theorem 1 easily. Let $U = U' \oplus \mathbf{0}$ (with $U' = (X^{t'})^{-1}$), then the matrix U is just the operation which Bob's performed on his qubit.

After introduced a new notation for the quantum teleportation, we obtained the condition on quantum teleportation by a generally entangled channel, the channel must not be maximally entangled state. We constructed the operation U performed by the Bob on the third qubit in order to obtain perfect replica of teleported state. We given the general

form of the operator U for the case that both channel and the state to be teleported are two-dimensional.

References

- [1] C. H. Bennett et al. Phys. Rev. Lett. **70** 1895 (1993).
- [2] L. Vaidman, Phys.Rev. A **49** 1473 (1994).
- [3] S. L. Braunstein and H. J. Kimble, Phys. Rev. Lett. **80** 869 (1998).
- [4] S. Bose and V. Vedral, Phys. Rev. A **61**, 040101(2000).
- [5] S. L. Braunstein, G. M. D'Ariano, G.L. Milburn and M. F. Sacchi, Phys.Rev.Lett. **84** 3486 (2000).
- [6] L. Accardi and M. Ohya, *quant-ph/9912087 (xxx.lanl.gov)*
- [7] Sergio Albeverio and Shao-Ming Fei, *quant-ph/0012035 (xxx.lanl.gov)*
- [8] Bei Zeng, Xiao Shu Liu, Yan Song Li and Gui Lu Long, *quant-ph/0104102 (xxx.lanl.gov)*
- [9] Wang-Li Li, Chuang-Feng Li and Guang-Can Guo, Phys. Rev. A **61**, 034301(2000)

